arXiv:1606.08752v2 [math.CV] 17 Nov 2016

ON QUADRATIC POLYNOMIAL MAPPINGS $f : \mathbb{C}^2 \to \mathbb{C}^2$

M. FARNIK & Z. JELONEK

Abstract. We show that up to linear equivalence, there is only finitely many polynomial quadratic mappings $f : \mathbb{C}^2 \to \mathbb{C}^2$ and $f : \mathbb{R}^2 \to \mathbb{R}^2$. We list all possibilities.

1. Introduction

Let $\Omega(d_1, d_2)$ denote the space of polynomial mappings $f = (f_1, f_2) : \mathbb{C}^2 \to \mathbb{C}^2$, where $\deg f_1 \leq d_1$ and $\deg f_2 \leq d_2$. Let $f, g \in \Omega(d_1, d_2)$. We say that f is topologically (respectively linearly) equivalent to g, if there are homeomorphisms (respectively linear isomorphisms) $\Phi, \Psi : \mathbb{C}^2 \to \mathbb{C}^2$, such that $f = \Psi \circ g \circ \Phi$. In the paper [1] it was showed that there is only a finite number of different topological types of mappings in $\Omega(d_1, d_2)$. Moreover, we know (see e.g. [5]) that there is a Zariski open dense subset $U \subset \Omega(d_1, d_2)$ such that every mapping $f \in U$ has the same, generic, topological type. If a mapping f has a generic topological type, then we say that f is a generic mapping. In practice it is difficult to describe the generic topological type and other topological types effectively.

Here we consider the case $d_1 = d_2 = 2$. We show that in this case the topological equivalence almost coincides with the linear equivalence. Moreover, we obtain a full classification of quadratic mappings of $\mathbb{C}^2$, with respect to the linear (hence also topological) equivalence. In particular we find a model of a generic mapping of $\Omega(2, 2)$.

We explain our basic idea. It was proved in [2] that a generic polynomial mapping from $\Omega(2, 2)$ has a rational cuspidial curve of degree 4 as a discriminant, moreover this curve is tangent in two smooth points to the line at infinity. On the other hand any two such rational cuspidial curves are projectively equivalent (see [7]), hence it is easy to deduce that discriminants of generic mappings from $\Omega(2, 2)$ are linearly equivalent. Using [3] we can deduce that any two generic mappings from $\Omega(2, 2)$ are algebraically equivalent and consequently (since they have the same algebraic degree) they are linearly equivalent. Moreover, there are only finitely many possible discriminants of quadratic mappings, up to a linear equivalence (because in the non-generic case they have a non-trivial action of an infinite affine group). Hence we can expect that there is only a finite number of orbits of the action of a linear group on $\Omega(2, 2)$. We show that it is indeed the case. In fact, it can be done in a very elementary way, however it is suprising that this result was not discovered up till now.

Let $C(f), \Delta(f), \mu(f)$ denote : the set of critical points, the discriminant and topological degree (see Definition 2.1). We have the following possibilities:

Generically-finite mappings:

(1) (the generic case) $f_1 = (x^2 + y, y^2 + x)$, $C(f_1) = \{4xy - 1 = 0\}$ is a hyperbola and $\Delta(f_1) = \{2^8x^2y^2 - 2^8x^3 - 2^8y^3 + 2^5 \cdot 9xy - 27 = 0\}$ is a reduced and irreducible curve with

Date: November 18, 2016.

1991 *Mathematics Subject Classification.* 14 D 99, 14 R 99, 51 M 99.

The authors were partially supported by the Narodowe Centrum Nauki grant number 2015/17/B/ST1/02637.

3 cusps at points $f_1(\frac{\varepsilon}{2}, \frac{\varepsilon^2}{2})$, where $\varepsilon^3 = 1$, $\dim O(f_1) = 12$, $O(f_1)$ is an open and dense affine subvariety of $\Omega(2,2)$, moreover $\chi(O(f_1)) = 0$ and $\mu(f_1) = 4$.

(2) $f_2 = (x^2 + y, xy)$, $C(f_2) = \{2x^2 = y\}$ is a parabola and $\Delta(f_2) = \{4x^3 = 27y^2\}$ is a cusp, $\dim O(f_2) = 11$, $O(f_2)$ is an affine subvariety of $\Omega(2,2)$, $\mu(f_2) = 3$.

(3) $f_3 = (x^2 + y, y^2)$, $C(f_3) = \{4xy = 0\}$ is two intersecting lines and $\Delta(f_3) = \{y(y - x^2) = 0\}$ is the sum of a line and a parabola, $\dim O(f_3) = 11$, $\mu(f_3) = 4$.

(4) $f_4 = (x^2, y^2)$, $C(f_4) = \{4xy = 0\}$ is two intersecting lines and $\Delta(f_4) = \{xy = 0\}$ is also two intersecting lines, $\dim O(f_4) = 10$ and $\mu(f_4) = 4$.

(5) $f_5 = (x^2 - x, xy)$, $C(f_5) = \{2x^2 - x = 0\}$ is two parallel lines and $\Delta(f_5)$ is the sum of the line $x = -1/4$ and the line $x = 0$, $\dim O(f_5) = 10$, $\mu(f_5) = 2$ and f_5 is not proper.

(6) $f_6 = (x^2, xy)$, $C(f_6) = \{x^2 = 0\}$ is a double line and $\Delta(f_6)$ is the line $x = 0$, $\dim O(f_6) = 9$, $\mu(f_6) = 2$ and f_6 is not proper.

(7) $f_7 = (xy, x + y)$, $C(f_7) = \{y = x\}$ is a line and $\Delta(f_7) = \{4x - y^2 = 0\}$ is a parabola, $\dim O(f_7) = 10$ and $\mu(f_7) = 2$.

(7') $f_9 = (x^2, y)$, $C(f_9) = \{x = 0\}$ is a line and $\Delta(f_9) = \{x = 0\}$ is also a line, $\dim O(f_9) = 9$ and $\mu(f_9) = 2$.

(8) $f_8 = (x, xy)$, $C(f_8) = \{x = 0\}$ is a line and $\Delta(f_8)$ is the line $\{x = 0\}$, $\dim O(f_8) = 9$, $\mu(f_8) = 1$ and f_8 is not proper.

(9) $f_{10} = (x^2 + y, x)$, $\dim O(f_{10}) = 8$, $C(f_{10})$ is the empty set and f_{10} is an automorphism, $\mu(f_{10}) = 1$.

(9') $f_{12} = (x, y)$, $\dim O(f_{12}) = 6$, $C(f_{12})$ is the empty set and f_{12} is an automorphism, $\mu(f_{12}) = 1$.

Not generically-finite mappings:

(10) $f_{11} = (x^2, x)$, $\dim O(f_{11}) = 7$ and $C(f_{11})$ is the plane.

(10') $f_{14} = (x^2 + y, 0)$, $\dim O(f_{14}) = 7$ and $C(f_{14})$ is the plane.

(10") $f_{16} = (x, 0)$, $\dim O(f_{16}) = 5$ and $C(f_{16})$ is the plane.

(11) $f_{13} = (xy, 0)$, $\dim O(f_{13}) = 8$ and $C(f_{13})$ is the plane.

(12) $f_{15} = (x^2, 0)$, $\dim O(f_{15}) = 6$ and $C(f_{15})$ is the plane.

(13) $f_{17} = (0, 0)$, $\dim O(f_{17}) = 2$ and $C(f_{17})$ is the plane.

Note that mappings f_7 and f_9 are topologically (even algebraically) equivalent. Similarly mappings f_{10}, f_{12} and f_{11}, f_{14}, f_{16} Hence we have 13 different topological types and 17 different linear types.

In Figure 1 we present the structure of $\Omega(2,2)$. Each row consists of orbits of given dimension, from the largest to the smallest, and a rising path joins two orbits if the smaller is contained in the closure of the larger.



FIGURE 1. The orbits

We also obtain similar results in the real case. We have the following possibilities (here $\Delta(f) = f(C(f))$):

Generically-finite mappings:

(1) (the first generic case) $f_1 = (x^2 + y, y^2 + x)$, $C(f_1) = \{4xy - 1 = 0\}$ is a hyperbola and $\Delta(f_1) = \{2^8x^2y^2 - 2^8x^3 - 2^8y^3 + 2^5 \cdot 9xy - 27 = 0\}$ is a reduced and irreducible curve with a cusp at $f_1(\frac{1}{2}, \frac{1}{2})$, $\dim O(f_1) = 12$, $O(f_1)$ is an open semi-algebraic subvariety of $\Omega(2, 2)$.

(1a) (the second generic case) $f_{1'} = \left(x^2 - y^2 + x, 2xy - y\right)$ with $C(f_{1'}) = \left\{4x^2 + 4y^2 - 1 = 0\right\}$ a circle and $\Delta(f_{1'}) = \{2^{16}(x^2 + y^2)^2 + 2^{17}(-x^3 + 3xy^2) + 2^9 \cdot 3^3 \cdot 5(x^2 + y^2) - (15)^3\}$ a reduced and irreducible curve with 3 cusps at points $f_{1'}\left(\frac{1}{4}, \frac{-\sqrt{3}}{4}\right)$, $f_{1'}\left(\frac{1}{4}, \frac{\sqrt{3}}{4}\right)$ and $f_{1'}\left(\frac{1}{2}, 0\right)$, $\dim O(f_{1'}) = 12$, $O(f_{1'})$ is an open semi-algebraic subvariety of $\Omega(2, 2)$.

(2) $f_2 = (x^2 + y, xy)$, $C(f_2) = \{2x^2 = y\}$ is a parabola and $\Delta(f_2) = \{4x^3 = 27y^2\}$ is a cusp, $\dim O(f_2) = 11$, $O(f_2)$ is a semi-algebraic subvariety of $\Omega(2, 2)$, $\mu(f_2) = 3$.

(3) $f_3 = (x^2 + y, y^2)$, $C(f_3) = \{4xy = 0\}$ is two intersecting lines and $\Delta(f_3) = \{y(y - x^2) = 0\}$ is the sum of a line and a parabola, $\dim O(f_3) = 11$.

the restriction of complex orbit of $f_4 = \left(x^2, y^2\right)$ splits into two orbits given by $\overline{\Phi}_1(f) < 0$ and $\overline{\Phi}_1(f) > 0$:

(4) the first case is represented by f_4 with $C(f_4) = \{4xy = 0\}$ two intersecting lines and $\Delta(f_4) = \{xy = 0\}$ also two intersecting lines, $\dim O(f_4) = 10$.

(4a) the second case is represented by $f_{4'} = \left(x^2 - y^2, xy\right)$ with $C(f_{4'}) = \{(0, 0)\}$ a point, $\Delta(f_{4'}) = \{(0, 0)\}$- a point, $\dim O(f_{4'}) = 10$.

(5) $f_5 = (x^2 - x, xy)$, $C(f_5) = \{2x^2 - x = 0\}$ is two parallel lines and $\Delta(f_5)$ is the sum of the line $x = -1/4$ and the point $(0, 0)$, $\dim O(f_5) = 10$, $\mu(f_5) = 2$ and f_5 is not proper.

(6) $f_6 = (x^2, xy)$, $C(f_6) = \{x^2 = 0\}$ is a double line and $\Delta(f_6)$ is the point $(0, 0)$, $\dim O(f_6) = 9$, $\mu(f_6) = 2$ and f_6 is not proper.

(7) $f_7 = (xy, x+y)$, $C(f_7) = \{y = x\}$ is a line and $\Delta(f_7) = \{4x - y^2 = 0\}$ is a parabola, $\dim O(f_7) = 10$ and $\mu(f_7) = 2$.

(7') $f_9 = (x^2, y)$, $C(f_9) = \{x = 0\}$ is a line and $\Delta(f_9) = \{x = 0\}$ is also a line, $\dim O(f_9) = 9$ and $\mu(f_9) = 2$.

(8) $f_8 = (x, xy)$, $C(f_8) = \{x = 0\}$ is a line and $\Delta(f_8)$ is the point $(0, 0)$, $\dim O(f_8) = 9$, $\mu(f_8) = 1$ and f_8 is not proper.

(9) $f_{10} = (x^2 + y, x)$, $\dim O(f_{10}) = 8$, $C(f_{10})$ is the empty set and f_{10} is an automorphism, $\mu(f_{10}) = 1$.

(9') $f_{12} = (x, y)$, $\dim O(f_{12}) = 6$, $C(f_{12})$ is the empty set and f_{12} is an automorphism, $\mu(f_{12}) = 1$.

Not generically-finite mappings:

(10) $f_{11} = (x^2, x)$, $\dim O(f_{11}) = 7$ and $C(f_{11})$ is the plane.

(10') $f_{14} = (x^2 + y, 0)$, $\dim O(f_{14}) = 7$ and $C(f_{14})$ is the plane.

(10") $f_{16} = (x, 0)$, $\dim O(f_{16}) = 5$ and $C(f_{16})$ is the plane.

(11) $f_{13} = (xy, 0)$, $\dim O(f_{13}) = 8$ and $C(f_{13})$ is the plane.

(12) $f_{15} = (x^2, 0)$, $\dim O(f_{15}) = 6$ and $C(f_{15})$ is the plane.

(13) $f_{17} = (0, 0)$, $\dim O(f_{17}) = 2$ and $C(f_{17})$ is the plane.

Thus we have 15 different topological real types and 19 different linear real types.

2. Main Result

Let us recall the following:

Definition 2.1. *If $f : \mathbb{C}^n \to \mathbb{C}^n$ is a generically-finite regular mapping, then $\Delta(f) = \{x \in \mathbb{C}^n : \#f^{-1}(x) \neq \mu(f)\}$ is called the discriminant of f. In particular, if f is proper, then $\Delta(f) = f(C(f))$, where $C(f)$ is the critical set of f. The set $\Delta(f)$ is either a hypersurface or the empty set (see e.g. [6]).*

We also use the following:

Definition 2.2. *By $GA(2,2)$ we denote the group of affine transformations of $\mathbb{C}^2$. By $\mathcal{G}A(2,2)$ we denote the group $GA(2,2) \times GA(2,2)$ with multiplication $(L_1, R_1) \circ (L_2, R_2) = (L_1 L_2, R_2 R_1)$. The group $\mathcal{G}A(2,2)$ acts on the set $\Omega(2,2)$: $(L, R)f = L \circ f \circ R$. We denote the orbit of $f \in \Omega(2,2)$ by $O(f)$. We say that $f_1, f_2 \in \Omega(2,2)$ are linearly equivalent, if there is a $\alpha \in \mathcal{G}A(2,2)$ such that $f_2 = \alpha f_1$, i.e. $f_2 \in O(f_1)$.*

We denote by $\mathrm{Stab}(f)$ the stabilizer of f. Note that $\dim \mathrm{Stab}(f) + \dim O(f) = \dim \mathcal{G}A(2,2) = 12$. Moreover if $(L, R) \in \mathrm{Stab}(f)$ and the image of f is not contained in a linear subspace of $\mathbb{C}^2$ then R uniquely determines L.

Let $f_1 = (x^2 + y, y^2 + x)$. We have:

Theorem 2.3. *Let $f \in \Omega(2,2)$. The following conditions are equivalent:*

1. *$C(f)$ is a hyperbola.*
2. *$O(f)$ is an open dense subset of $\Omega(2,2)$.*
3. *$O(f) = O(f_1)$, i.e. f is linearly equivalent to f_1.*

Moreover, the maximal orbit $O(f_1)$ is a smooth affine variety of Euler characteristic 0 and dimension 12.

Proof. (1) $\Rightarrow$ (2) First note that either the mapping $f|_{C(f)}$ has a critical point or $\Delta(f)$ is a singular curve. Indeed, in other case $\Delta(f)$ is isomorphic to a hyperbola and the curve $f^{-1}(\Delta(f))$ has the Euler characteristic equal to 0 (as a covering of hyperbola). In particular $\mu(f)\chi(\mathbb{C}^2 \setminus \Delta(f)) = \chi(\mathbb{C}^2 \setminus f^{-1}(\Delta(f))) = 1$ and $\mu(f) = 1$, a contradiction.

Now we will show that $\mathrm{Stab}(f)$ is finite. If $(L, R) \in \mathrm{Stab}(f)$ then R must preserve $C(f)$, all singular points of $f|_{C(f)}$ and all pre-images of singular points of $\Delta(f)$. Hence R has to preserve a hyperbola and some non-empty finite subset of the hyperbola. It is easy to see that there is only a finite number of such linear mappings R.

Since $\mathrm{Stab}(f)$ is finite we know that $\dim O(f) = \dim \Omega(2,2)$ and $O(f)$ is a maximal orbit, in particular it is open and dense.

(2) $\Rightarrow$ (3) It is easy to check that $C(f_1)$ is a hyperbola, hence $\mathcal{G}A(2,2)f_1$ is open and dense. In particular $O(f) \cap O(f_1) \neq \emptyset$.

(3) $\Rightarrow$ (1) Obvious.

Now we will describe the set $\mathrm{Stab}(f_1)$. Let us note that $C(f_1) = \{(x, y) : 4xy = 1\}$. Hence f_1 has exactly three $A_{1,1}$ singularities $P_i = (-1/2\alpha, -1/2\alpha^2)$, where $\alpha^3 = 1$. Note that the linear automorphisms which preserve $C(f_1)$ and P_i for $i = 1, 2, 3$ are exactly the mappings $(x, y), (\alpha x, \alpha^2 y), (\alpha^2 x, \alpha y)$ and $(y, x), (\alpha y, \alpha^2 x), (\alpha^2 y, \alpha x)$. Hence $\mathrm{Stab}(f_1)$ has exactly 6 elements. In order to calculate the Euler characteristic of the maximal orbit $O = \mathcal{G}A(2,2)f_1$ note that the mapping $\Psi \colon \mathcal{G}A(2,2) \ni (L, R) \mapsto L \circ f_1 \circ R \in O$ is a topological covering of degree 6. Hence $\chi(\mathcal{G}A(2,2)) = 6\chi(O)$. Since $GA(2,2)$ is homotopic to $\mathbb{C}^4$ minus a cone, it has Euler characteristic equal to 0. Since $\mathcal{G}A(2,2) = GA(2,2) \times GA(2,2)$ we have $\chi(\mathcal{G}A(2,2)) = 0$. Hence $\chi(O) = 0$. □

Remark 2.4. We say that $f \in \Omega(2,2)$ is general if $O(f)$ is dense in $\Omega(2,2)$. Hence a general mapping is equivalent to $(x^2 + y, y^2 + x)$. Now assume that $f \in \Omega(2,2)$ is not a general mapping. From the proof of Theorem 2.3 we conclude that there are infinite algebraic groups of affine mappings of $\mathbb{C}^2$, which preserve $C(f)$, $\Delta(f)$ and $f^{-1}(\Delta(f))$. Hence the curves $C(f)$, $\Delta(f)$, $f^{-1}(\Delta(f))$ are very special. In particular, the curve $f^{-1}(\Delta(f))$ is rational.

Remark 2.5. From our classification we have that a generic mapping is general. In other words a generic mapping is linearly equivalent to $(x^2 + y, y^2 + x)$.

Now we will describe the other orbits. First we will show that there are exactly two orbits of dimension 11. Let $f \in \Omega(2,2)$, we can write $f = (g, h)$ where

$$g = a_1x^2 + b_1xy + c_1y^2 + d_1x + e_1y + f_1,$$

$$h = a_2x^2 + b_2xy + c_2y^2 + d_2x + e_2y + f_2.$$

Let $A = 2a_1b_2 - 2a_2b_1$, $B = 4a_1c_2 - 4a_2c_1$, $C = 2b_1c_2 - 2b_2c_1$, $D = 2a_1e_2 + d_1b_2 - 2a_2e_1 - d_2b_1$, $E = 2d_1c_2 + b_1e_2 - 2d_2c_1 - b_2e_1$ and $F = d_1e_2 - d_2e_1$, the Jacobian of f is given by the formula:

$$J(f) = Ax^2 + Bxy + Cy^2 + Dx + Ey + F.$$

Consider the following two matrices:

$$\Phi_1(f) = \begin{bmatrix} 2A & B \\ B & 2C \end{bmatrix} \quad \text{and} \quad \Phi_2(f) = \begin{bmatrix} 2A & B & D \\ B & 2C & E \\ D & E & 2F \end{bmatrix}$$

Note that polynomial $\overline{\Phi}_1 := \det \Phi_1$ is a quadratic function of c_1 and $\overline{\Phi}_2 := \det \Phi_2$ is a quadratic function of d_1. Using the obvious lemma below it is easy to check that the polynomials $\overline{\Phi}_1$ and $\overline{\Phi}_2$ are irreducible.

Lemma 2.6. *Let* $\Phi = a(x)y^2+b(x)y+c(x) \in \mathbb{C}[x_1, \dots, x_n, y]$, *where* $\gcd(a(x), b(x), c(x)) = 1$. *Then* Φ *is reducible if and only if* $b^2 - 4ac$ *is a square in* $\mathbb{C}[x_1, \dots, x_n]$.

Obviously the critical set $C(f)$ is a quadric if and only if $\operatorname{rank} \Phi_1 > 0$. In this case $\overline{\Phi}_1(f) = 0$ if and only if $C(f)$ has only one point at infinity, moreover $\overline{\Phi}_2(f) = 0$ if and only if the projective curve $\overline{C(f)}$ is singular, i.e. $C(f)$ is a sum of two lines, finally, $\operatorname{rank} \Phi_2(f) = 1$ if and only if $C(f)$ is a double line. In particular $O(f_1) = \Omega(2,2) \setminus (\{\overline{\Phi}_1 = 0\} \cup \{\overline{\Phi}_2 = 0\})$, which confirms in a different way that $O(f_1)$ is an affine variety. Now we will describe the set $\{\overline{\Phi}_1 = 0\} \setminus \{\overline{\Phi}_2 = 0\}$. Let $f_2 = (x^2 + y, xy)$, we have:

Theorem 2.7. *The following conditions are equivalent:*

(1) $O(f) = \{\overline{\Phi}_1 = 0\} \setminus \{\overline{\Phi}_2 = 0\}$.
(2) $C(f)$ *is a parabola.*
(3) $O(f) = O(f_2)$, *i.e.* f *is linearly equivalent to* f_2.

In particular the orbit $O(f_2)$ *is a smooth affine variety of dimension* 11.

Proof. (1) $\Rightarrow$ (2) It is obvious.

(2) $\Rightarrow$ (3) First observe that $f = (g, h)$ is a proper mapping. Indeed, if f is not proper, then by [4] the curve $C(f)$ must be contracted by f to some point (a, b). Hence $g-a, h-b$ are proportional to the equation of $C(f)$ and $J(f)$ vanishes on $\mathbb{C}^2$, a contradiction.

We claim that $\Delta(f)$ is not smooth. In the other case $\Delta(f) \cong \mathbb{C}$ and by [3] the mapping f is algebraically equivalent to (x^r, y). Hence f must be linearly equivalent to (x^2, y) and $C(f)$ is not a parabola, a contradiction. Consequently the curve $\Delta(f)$ has a singular point. This implies that the group of linear mappings which stabilize $\Delta(f)$ has dimension at most 1. On the other hand f is not generic so the group $\operatorname{Stab}(f)$ in $\mathcal{G}A(2,2)$ has dimension exactly one. Hence $\dim O(f) = 11 = \dim\{\overline{\Phi}_1 = 0\}$. This means that $O(f)$ is open and dense in $\{\overline{\Phi}_1 = 0\}$. By the same reason $O(f_2)$ is open and dense in $\{\overline{\Phi}_1 = 0\}$. Since the hypersurface $\{\overline{\Phi}_1 = 0\}$ is irreducible, we have $O(f) \cap O(f_2) \neq \emptyset$. Hence $O(f_2) = O(f)$.

(3) $\Rightarrow$ (1) It is obvious. □

Remark 2.8. Note that the mapping f_2 is a pattern of a cusp singularity.

Our next aim is to describe the set $\{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}$. Let $f_3 = (x^2 + y, y^2)$. We have:

Theorem 2.9. *The following conditions are equivalent:*

(1) $C(f)$ *is isomorphic to a cross* $K_2 = \{xy = 0\}$ *and* $\Delta(f)$ *contains a parabola.*
(2) $O(f)$ *is dense in* $\{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}$.
(3) $O(f) = O(f_3)$, *i.e.* f *is linearly equivalent to* f_3.

In particular the orbit $O(f_3)$ *is a smooth variety of dimension* 11.

Proof. (1) $\Rightarrow$ (2) Note that f is a proper mapping. If $f = (g, h)$ then $f = 0$ and $g = 0$ have no common points at infinity. Indeed, we can assume that $C(f) = K_2$. Hence the common zeroes of g and h at the line at infinity are of the form $(1 : 0)$ or $(0 : 1)$. By symmetry it is enough to consider the first case. Hence $a_1 = a_2 = 0$ and consequently $B = 0$. This implies that $J(f) \neq xy$. This contradiction shows that the curves $g = a$,$h = b$ have no common points at infinity, hence $\mu(f) = 4$ and f is proper.

Note that $\Delta(f) = P \cup P_1$, where P is a parabola and $P_1 \neq P$ is a line or a parabola. Indeed, we can assume that $P = \{y = x^2\}$. Now, if $\Delta(f) = P$ then by [3] f is algebraically equivalent to (x^r, y), hence $C(f)$ is not isomorphic to K_2, a contradiction. Denote $R = P \cap P_1$, we can assume that $R = (0, 0)$.

Consider the group $\mathrm{Stab}(f) \subset \mathcal{G}A(2, 2)$. Every affine transformation which preserves (P, R) is of the form (ax, a^2y). Hence $\dim \mathrm{Stab}(f) \leq 1$ and since f is not generic we have $\dim \mathrm{Stab}(f) = 1$. Consequently $\dim O(f) = 11$ and $O(f)$ is dense in $\{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}$.

(2) $\Rightarrow$ (3) It is obvious.

(3) $\Rightarrow$ (1) It is obvious. □

Remark 2.10. It is easy to see that the discriminant of f_3 (and hence the discriminant of every mapping from $O(f_3)$) is a union of a parabola and a line which is tangent to this parabola.

Let $f_4 = (x^2, y^2)$. We will describe the set $(\{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}) \setminus O(f_2)$.

Theorem 2.11. *The following conditions are equivalent:*

(1) *$C(f)$ and $\Delta(f)$ are isomorphic to the cross $K_2 = \{xy = 0\}$.*
(2) *$O(f) = O(f_4)$, i.e. f is linearly equivalent to f_4.*
(3) *$O(f) = (\{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}) \setminus O(f_3)$.*

The orbit $O(f_4)$ is a smooth variety of dimension 10. *Moreover, $O(f_3) \cup O(f_4) = \{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}$.*

Proof. (1) $\Rightarrow$ (2) We can assume that $C(f) = K_2$ and $\Delta(f) = K_2$. As in the proof above, the mapping f is proper. Hence by [3] the mapping f is algebraically equivalent to (x^r, y^s) for some $r, s > 1$. Since $\mu(f) \leq 4$ we have $r = s = 2$. Finally f is linearly equivalent to (x^2, y^2) because $f \in \Omega(2, 2)$.

(2) $\Rightarrow$ (3) If $f \in \{\overline{\Phi}_2 = 0\} \setminus \{\overline{\Phi}_1 = 0\}$ and $O(f) \neq O(f_3)$, then $\Delta(f)$ is isomorphic either to K_2 or to a line. The second possibility can be excluded. Indeed, assume that $\Delta(f)$ is a line. By [3] the mapping f is algebraically equivalent to a mapping (x^r, y) hence $C(f)$ is not isomorphic to K_2, a contradiction.

(3) $\Rightarrow$ (1) We have $f_4 \in O(f)$.

Finally $\mathrm{Stab}(f_4) = \{((\alpha^{-2}x, \beta^{-2}y), (\alpha x, \beta y)), \alpha, \beta \in \mathbb{C}^*\} \cup \{((\beta^{-2}y, \alpha^{-2}x), (\alpha y, \beta x)), \alpha, \beta \in \mathbb{C}^*\}$, hence $\dim \mathrm{Stab}(f_4) = 2$ and consequently $\dim O(f_4) = 10$. □

Now we would like to obtain the equations of the closure of $O(f_4)$. Note that if $f = (g, h) = f_4 \circ R$, where $R \in GA(2, 2)$ then g and h are squares, i.e. the coefficients of g and h satisfy:

$$\mathrm{rank} \begin{bmatrix} 2a_1 & b_1 & d_1 \\ b_1 & 2c_1 & e_1 \\ d_1 & e_1 & 2f_1 \end{bmatrix} \leq 1 \quad \text{and} \quad \mathrm{rank} \begin{bmatrix} 2a_2 & b_2 & d_2 \\ b_2 & 2c_2 & e_2 \\ d_2 & e_2 & 2f_2 \end{bmatrix} \leq 1$$

Furthermore if we take $f = (g, h) = L \circ f_4 \circ R$, where $(L, R) \in \mathcal{G}A(2, 2)$ then $g - f_1$ and $h - f_2$ are linear combinations of two squares (both of the same two). Thus the coefficients of f satisfy the condition $\operatorname{rank} \Psi_1 \leq 2$, where:

$$\Psi_1 = \begin{bmatrix} 2a_1 & b_1 & 2a_2 & b_2 \\ b_1 & 2c_1 & b_2 & 2c_2 \\ d_1 & e_1 & d_2 & e_2 \end{bmatrix}.$$

Obviously $O(f_4)$ is a dense open subset of the set $\{\operatorname{rank} \Psi_1 \leq 2\}$. In particular we obtain that $O(f_3) = \{\overline{\Phi}_2 = 0\} \backslash (\{\overline{\Phi}_1 = 0\} \cup \{\operatorname{rank} \Psi_1 \leq 2\})$ and $O(f_4) = \{\operatorname{rank} \Psi_1 \leq 2\} \backslash \{\overline{\Phi}_1 = 0\}$.

The remaining part of orbits is contained in $\{\overline{\Phi}_2 = 0\} \cap \{\overline{\Phi}_1 = 0\}$, in particular all of them have codimension greater than or equal to 2. The rest of the paper is devoted to description of these orbits. Note that if $f \in \{\overline{\Phi}_2 = 0\} \cap \{\overline{\Phi}_1 = 0\}$, then $C(f)$ is either two parallel lines or a double line or a line or it is the empty set or the whole $\mathbb{C}^2$.

Let $f_5 = (x^2 - x, xy)$. We will describe the set $\{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 2\}$.

Theorem 2.12. *The following conditions are equivalent:*

(1) $O(f) = \{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 2\}$.
(2) $C(f)$ *consists of two parallel lines.*
(3) $O(f) = O(f_5)$, *i.e.* f *is linearly equivalent to* f_5.

The orbit $O(f_5)$ *is a smooth variety of dimension* 10.

Proof. (1) $\Rightarrow$ (2) Obvious.

(2) $\Rightarrow$ (3) By composing with suitable R we may assume that $J(f) = 2x^2 - x$, hence $2 = A = 2a_1b_2 - 2a_2b_1$. So by composing with L we may additionally assume that $a_1 = 1$ and $a_2 = 0$, hence $b_2 = 1$ and we may assume that $b_1 = 0$. Furthermore $0 = B = 4c_2$, $0 = C = -2c_1$, $-1 = D = 2e_2 + d_1$, $0 = E = -e_1$ and $0 = F = d_1e_2$. Hence either $f = (x^2 + f_1, xy + d_2x - \frac{1}{2}y + f_2)$ or $f = (x^2 - x, xy + d_2x + f_2)$. In both cases f is equivalent to f_5.

(3) $\Rightarrow$ (1) In virtue of the implication (2) $\Rightarrow$ (3) it is obvious.

Finally the set $\{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 2\}$ is an open subset of $\{\overline{\Phi}_1 = 0\} \cap \{\overline{\Phi}_2 = 0\}$. Since the latter set has dimension 10 we have $\dim O(f_5) = 10$. □

Let $f_6 = (x^2, xy)$. We will describe the set $\{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 1\}$.

Theorem 2.13. *The following conditions are equivalent:*

(1) $C(f)$ *is a double line.*
(2) $O(f) = O(f_6)$, *i.e.* f *is linearly equivalent to* f_6.
(3) $O(f) = \{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 1\}$.

The orbit $O(f_6)$ *is a smooth variety of dimension* 9.

Proof. (1) $\Rightarrow$ (2) As in Theorem 2.12 we can assume that $J(f) = 2x^2$, $a_1 = b_1 = 1$ and $a_2 = b_2 = 0$. Then it follows that $c_1 = c_2 = d_1 = e_1 = e_2 = 0$. Hence $f = (x^2 + f_1, xy + d_2x + f_2)$ and is equivalent to f_6.

(2) $\Rightarrow$ (3) Note that for every $f \in \{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 1\}$ the set $C(f)$ is a double line. Hence by the first part of the proof, we have $\{\operatorname{rank} \Phi_1 = 1\} \cap \{\operatorname{rank} \Phi_2 = 1\} = O(f_6)$.

(3) $\Rightarrow$ (1) Obvious.

Finally note that $\operatorname{Stab}(f_6) = \{(\alpha^{-2}x, (\alpha\gamma)^{-1}y - \beta\gamma^{-1}\alpha^{-2}x), (\alpha x, \beta x + \gamma y), \alpha, \gamma \in \mathbb{C}^*, \beta \in \mathbb{C}\}$. Hence we have $\dim O(f_6) = 9$. □

From the theorems above it follows that $\{\operatorname{rank}\Phi_1 > 0\} = O(f_1) \cup \ldots \cup O(f_6)$. We will now focus on the mappings satisfying $\operatorname{rank}\Phi_1(f) = 0$. Let

$$\Phi_3(f) = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{bmatrix} \quad \text{and} \quad \Phi_4(f) = \begin{bmatrix} a_1 & b_1 & c_1 & d_1 & e_1 \\ a_2 & b_2 & c_2 & d_2 & e_2 \end{bmatrix}.$$

The assumption $\operatorname{rank}\Phi_1(f) = 0$ is equivalent to $\operatorname{rank}\Phi_3(f) < 2$ and to $C(f)$ not being a quadric. In particular $O(f) \cap \Omega(2,1) \neq \emptyset$, so from now on we will assume that $f \in \Omega(2,1)$, i.e. $a_2 = b_2 = c_2 = 0$.

Let $f = (g_2 + g_1 + g_0, h_1 + h_0)$, where g_i and h_i are homogeneous of degree i. We may assume that $\deg g \geq \deg h$ then $\operatorname{rank}\Phi_3(f) = 0$ if and only if $g_2 = 0$, $\operatorname{rank}\Phi_4(f) < 2$ if and only if $h_1 = 0$ and $\operatorname{rank}\Phi_4(f) = 0$ if and only if f is constant.

Now we will describe the case when $g_2 h_1 \neq 0$. Let $f_7 = (xy, x+y)$, $f_8 = (x, xy)$, $f_9 = (x^2, y)$, $f_{10} = (x^2 + y, x)$ and $f_{11} = (x^2, x)$.

Theorem 2.14. *Let $f = (g_2 + g_1 + g_0, h_1 + h_0)$ and $g_2 h_1 \neq 0$, then one of the following holds:*

1. *g_2 is not a square, h_1 does not divide g_2. Then $f \in O(f_7)$. Moreover $C(f)$ is a line, $\Delta(f)$ is a parabola and $\dim \operatorname{Stab}(f) = 2$. The mapping f is proper.*
2. *g_2 is not a square, h_1 divides g_2. Then $f \in O(f_8)$. Moreover $C(f)$ is a line and $\Delta(f)$ is a line and $\dim \operatorname{Stab}(f) = 3$. The mapping f is not proper.*
3. *g_2 is a square, h_1 does not divide g_2. Then $f \in O(f_9)$. Moreover $C(f)$ and $\Delta(f)$ are lines and $\dim \operatorname{Stab}(f) = 3$. The mapping f is proper.*
4. *g_2 is a square, h_1 divides g_2 but not g_1. Then $f \in O(f_{10})$. Moreover f is an automorphism and $\dim \operatorname{Stab}(f) = 4$. The mapping f is proper.*
5. *g_2 is a square, h_1 divides g_2 and g_1. Then $f \in O(f_{11})$. Moreover $C(f) = \mathbb{C}^2$, $\operatorname{Im}(f)$ is a parabola and $\dim \operatorname{Stab}(f) = 5$. The mapping f is not dominant.*

Proof. (1) Since g_2 is not a square it is a product of two independent linear forms and we may assume, by composing with R, that $g_2 = xy$. Moreover since h_1 does not divide g_2 we have $h_1 = d_2 x + e_2 y$ for $d_2, e_2 \in \mathbb{C}^*$. Composing with $R(x,y) = (x/d_2, y/e_2)$ and $L(x,y) = (d_2 e_2 x, y)$ we obtain $h_1 = x + y$ and retain $g_2 = xy$. Thus $f = (xy + d_1 x + e_1 y + f_1, x + y + f_2)$ and for $R(x,y) = (x - e_1, y - d_1)$ and $L(x,y) = (x + e_1 d_1 - f_1, y + d_1 + e_1 - f_2)$ we obtain $f_7 = L \circ f \circ R$. Clearly $C(f_7) = \{y = x\}$ is a line and $\Delta(f_7) = \{4x = y^2\}$ is a parabola. It remains to calculate $\dim \operatorname{Stab}(f_7)$. Since for every $(R, L) \in \operatorname{Stab}(f_7)$ the mapping R must preserve the parabola $\Delta(f)$ we see that $\dim \operatorname{Stab}(f_7) \leq 2$. On the other hand the codimension of $O(f_7)$ is a least two. Consequently $\dim \operatorname{Stab}(f_7) = 2$.

(2) Similarly as in (1) we may assume that $g_2 = xy$ and $h_1 = x$. It follows that f is equivalent with f_8. Moreover $J(f_8) = x$ and $\Delta(f_8) = \{x = 0\}$. Let $(L, R) \in \operatorname{Stab}(f_8)$ and $R = (R_1, R_2)$. Since R preserves $C(f_8)$ we have $R_1 = \alpha x$ and $R_2 = \beta y + \gamma$, where $\alpha, \beta \in \mathbb{C}^*$ and $\gamma \in \mathbb{C}$. Thus $f_8 \circ R(x,y) = (\alpha\beta xy + \alpha\gamma, \alpha x)$ and for $L(x,y) = ((x - \alpha\gamma)/\alpha\beta, y/\alpha)$ we obtain $L \circ f_8 \circ R = f_8$.

(3) We may assume that $g_2 = x^2$ and $h_1 = y$. It follows that f is equivalent with f_9. Moreover $J(f_9) = 2x$ and $\Delta(f_9) = \{x = 0\}$. As above, for $(L, R) \in \operatorname{Stab}(f_9)$ and $R = (R_1, R_2)$ we obtain $R_1 = \alpha x$ and $R_2 = \beta y + \gamma$, where $\alpha, \beta \in \mathbb{C}^*$ and $\gamma \in \mathbb{C}$.

(4) We may assume that $g_2 = x^2$, $g_1 = y$ and $h_1 = x$. It follows that f is equivalent with f_{10}. Let $(L, R) \in \operatorname{Stab}(f_{10})$ and $R = (R_1, R_2)$. Since R_1^2 is the only term in $f_{10} \circ R$ with quadratic part we have $R_1 = \alpha x + \beta$, where $\alpha \in \mathbb{C}^*$ and $\beta \in \mathbb{C}$. Moreover the coefficients at x^2 and y in the first component of $f_{10} \circ R$ must be equal, thus $R_2 = \gamma x + \alpha^2 y + \delta$, where $\gamma, \delta \in \mathbb{C}$.

(5) Similarly as in (1) we may assume that $g_2 = x^2$, $g_1 = x$ and $h_1 = x$. It follows that f is equivalent with f_{11}. Let $(L, R) \in \mathrm{Stab}(f_{11})$ and $R = (R_1, R_2)$. As in (4) we have $R_1 = \alpha x + \beta$, where $\alpha \in \mathbb{C}^*$ and $\beta \in \mathbb{C}$, however R_2 is arbitrary. □

Note that the conditions "g_2 is a square", "h_1 divides g_2" and "h_1 divides g_1" can be expressed in terms of the coefficients of $f \in \Omega(2,2)$. We will do it in the affine chart $a_1 \neq 0$ – one of the six natural charts covering the set $\{g_2 \neq 0\}$.

Obviously g_2 is a square if and only if $4a_1c_1 - b_1^2 = 0$. Moreover $h_1 = d_2x + e_2y - \frac{a_2}{a_1}(d_1x + e_1y)$, hence it divides g_1 if and only if $a_1(a_1e_2 - a_2e_1)^2 - b_1(a_1e_2 - a_2e_1)(a_1d_2 - a_2d_1) + c_1(a_1d_2 - a_2d_1)^2 = 0$. Finally, h_1 divides g_1 if and only if $d_1(a_1e_2 - a_2e_1) - e_1(a_1d_2 - a_2d_1) = 0$.

Notice that from $g_2 = 0$ and $h_1 \neq 0$ follows that $f \in GA(2,2)$. In this case $f \in O(f_{12})$, where $f_{12} = (x, y)$, and $\dim O(f) = 6$.

The last case to describe is $h_1 = 0$. Let $f_{13} = (xy, 0)$, $f_{14} = (x^2 + y, 0)$, $f_{15} = (x^2, 0)$, $f_{16} = (x, 0)$ and $f_{17} = (0, 0)$.

Theorem 2.15. *Let $f = (g_2 + g_1 + g_0, 0)$, then $C(f) = \mathbb{C}^2$ and one of the following holds:*

1. *g_2 is not a square. Then $f \in O(f_{13})$ and $\dim \mathrm{Stab}(f) = 4$.*
2. *g_2 is a square, g_1 is nonzero and does not divide g_2. Then $f \in O(f_{14})$ and $\dim \mathrm{Stab}(f) = 5$.*
3. *g_2 is a square, g_1 is zero or divides g_2. Then $f \in O(f_{15})$ and $\dim \mathrm{Stab}(f) = 6$.*
4. *$g_2 = 0$ and $g_1 \neq 0$. Then $f \in O(f_{16})$ and $\dim \mathrm{Stab}(f) = 7$.*
5. *$g = 0$. Then $f \in O(f_{17})$ and $\dim \mathrm{Stab}(f) = 10$.*

Proof. Note that in cases (1)–(4) the image of f is a line. Thus if $(L, R) \in \mathrm{Stab}(f)$ then R determines L up to an automorphism preserving every point of the line, the group of such automorphisms is two dimensional. In case (5) the image of f is a point, which is preserved by a four dimensional group.

(1) Since g_2 is not a square we may assume, by composing with R, that $g_2 = xy$. Thus $f = (xy + d_1x + e_1y + f_1, 0)$, which is equivalent with f_{13}. Let $(L, R) \in \mathrm{Stab}(f_{13})$ and $R = (R_1, R_2)$. We have $R_1R_2 = \alpha_1xy + \alpha_2$, consequently R is equal either to (β_1x, β_2y) or to (β_1y, β_2x), where $\beta_1, \beta_2 \in \mathbb{C}^*$.

(2) We may assume that $g_2 = x^2$ and $g_1 = y$, so f is equivalent with f_{14}. Let $(L, R) \in \mathrm{Stab}(f_{14})$ and $R = (R_1, R_2)$. The quadratic part of R_1^2 is equal $\alpha_1^2x^2$, thus $R = (\alpha_1x + \alpha_2, \alpha_1^2y - 2\alpha_1\alpha_2x + \alpha_3)$, where $\alpha_1 \in \mathbb{C}^*$ and $\alpha_2, \alpha_3 \in \mathbb{C}$.

(3) We may assume that $f = (x^2 + d_1x + f_1, 0)$, which is equivalent with f_{15}. Let $(L, R) \in \mathrm{Stab}(f_{15})$ and $R = (R_1, R_2)$. We have $R_1 = \alpha_x$ and R_2 arbitrary.

(4) and (5) are obvious.

□

3. The real case

In this section we will determine the orbits in $\Omega(2,2)$ over the field of real numbers. We will base on the complex case and to avoid confusion we will use the subscripts $\mathbb{R}$ and $\mathbb{C}$ to indicate over which field an object is considered. Obviously for $f \in \Omega_{\mathbb{R}}(2,2)$ we have $O_{\mathbb{R}}(f) \subset O_{\mathbb{C}}(f) \cap \Omega_{\mathbb{R}}(2,2)$. In most cases equality holds, however there are two cases when the restriction of a complex orbit splits into two real orbits. We have the following possibilities:

(1) the restriction of the generic complex orbit splits into two semialgebraic orbits given by $\overline{\Phi}_1(f) < 0$ and $\overline{\Phi}_1(f) > 0$. The first case is represented by $f_1 = \left(x^2 + y, y^2 + x\right)$ with $C(f_1) = \{4xy - 1 = 0\}$ a hyperbola and $\Delta(f_1) = \left\{2^8x^2y^2 - 2^8x^3 - 2^8y^3 + 2^5 \cdot 9xy - 27 = 0\right\}$ a reduced and irreducible curve with a cusp at $f_1\left(\frac{1}{2}, \frac{1}{2}\right)$. The second case is represented by $f_{1'} = \left(x^2 - y^2 + x, 2xy - y\right)$ with $C(f_{1'}) = \left\{4x^2 + 4y^2 - 1 = 0\right\}$ a circle and $\Delta(f_{1'}) = \{2^{16}(x^2 + y^2)^2 + 2^{17}(-x^3 + 3xy^2) + 2^9 \cdot 3^3 \cdot 5(x^2 + y^2) - (15)^3\}$ a reduced and irreducible curve with 3 cusps at points $f_{1'}\left(\frac{1}{4}, \frac{-\sqrt{3}}{4}\right)$, $f_{1'}\left(\frac{1}{4}, \frac{\sqrt{3}}{4}\right)$ and $f_{1'}\left(\frac{1}{2}, 0\right)$.

(2) $O_{\mathbb{R}}(f_2) = O_{\mathbb{C}}(f_2) \cap \Omega_{\mathbb{R}}(2, 2)$.

(3) $O_{\mathbb{R}}(f_3) = O_{\mathbb{C}}(f_3) \cap \Omega_{\mathbb{R}}(2, 2)$.

(4) the restriction of complex orbit of $f_4 = \left(x^2, y^2\right)$ splits into two orbits given by $\overline{\Phi}_1(f) < 0$ and $\overline{\Phi}_1(f) > 0$. The first case is represented by f_4 with $C(f_4) = \{4xy = 0\}$ two intersecting lines and $\Delta(f_4) = \{xy = 0\}$ also two intersecting lines. The second case is represented by $f_{4'} = \left(x^2 - y^2, xy\right)$ with $C(f_{1'}) = \left\{2x^2 + 2y^2 = 0\right\}$ a point.

(5) for $f \in \{f_5, \ldots, f_{17}\}$ we have $O_{\mathbb{R}}(f) = O_{\mathbb{C}}(f) \cap \Omega_{\mathbb{R}}(2, 2)$.

Proof. (1) Let $f \in O_{\mathbb{C}}(f_1) \cap \Omega_{\mathbb{R}}(2, 2)$. Note that f has 3 cusps. Since the cusps are given by real equations they are either real or in complex conjugate pairs. So f may have either 3 or 1 real cusp. Note that $f_{1'} = \left(x^2 - y^2 + x, 2xy - y\right)$ has 3 real cusps and f_1 has 1.

Assume that f has 3 real cusps. There are some $L, R \in GA_{\mathbb{C}}(2, 2)$ such that $f_{1'} = L \circ f \circ R$. Note that R maps the three real and non-collinear cusps of $f_{1'}$ into the three real cusps of f. Hence $R(\mathbb{R}^2) = \mathbb{R}^2$, so $R \in GA_{\mathbb{R}}(2, 2)$. Furthermore $f \circ R(\mathbb{R}^2) = \mathbb{R}^2$, hence also $L \in GA_{\mathbb{R}}(2, 2)$. Thus $f \in O_{\mathbb{R}}(f_{1'})$.

Now assume that f has 1 real cusp. There are some $L, R \in GA_{\mathbb{C}}(2, 2)$ such that $f_1 = L \circ f \circ R$. By composing with an element of $\mathrm{Stab}(f_1)$ (see the proof of Theorem 2.3) we may assume that R maps the real cusp of f_1 to the real cusp of f. Since the complex cusps are conjugate the mapping $\overline{R}$, i.e. the conjugate of R, coincides with R on the three non-collinear cusps. Hence $\overline{R} = R$, so $R \in GA_{\mathbb{R}}(2, 2)$ and consequently $L \in GA_{\mathbb{R}}(2, 2)$ and $f \in O_{\mathbb{R}}(f_1)$.

Now we will show that if $f' \in O_{\mathbb{R}}(f)$ then $\mathrm{sgn}(\overline{\Phi}_1(f')) = \mathrm{sgn}(\overline{\Phi}_1(f))$. Indeed, let $L, R \in GA_{\mathbb{R}}(2, 2)$, we have $J(L \circ f \circ R) = (J(L) \circ f \circ R) \cdot (J(f) \circ R) \cdot J(R) = J(L) \cdot J(R) \cdot (J(f) \circ R)$. Note that $\Phi_1(f)$ is the matrix defining the quadratic form which is the homogeneous part of $2J(f)$ of degree 2. Consequently $\Phi_1(L \circ f \circ R) = J(L) \cdot J(R) \cdot D(R)^T \cdot \Phi_1(f) \cdot D(R)$, where $D(R)$ is the derivative of the affine map R. Hence $\overline{\Phi}_1(L \circ f \circ R) = J(L)^2 \cdot J(R)^2 \cdot \det(D(R)^T) \cdot \det(\Phi_1(f)) \cdot \det(D(R)) = J(L)^2 \cdot J(R)^4 \cdot \overline{\Phi}_1(f)$. So in particular $\mathrm{sgn}(\overline{\Phi}_1(L \circ f \circ R)) = \mathrm{sgn}(\overline{\Phi}_1(f))$.

As a consequence we obtain that $f \in O_{\mathbb{R}}(f_{1'})$ if and only if $\overline{\Phi}_1(f) > 0$ and $f \in O_{\mathbb{R}}(f_1)$ if and only if $\overline{\Phi}_1(f) < 0$. Indeed if $f \in O_{\mathbb{C}}(f_1) \cap \Omega_{\mathbb{R}}(2, 2)$ then either $f \in O_{\mathbb{R}}(f_1)$ and $\mathrm{sgn}(\overline{\Phi}_1(f)) = \mathrm{sgn}(\overline{\Phi}_1(f_1)) = -1$ or $f \in O_{\mathbb{R}}(f_{1'})$ and $\mathrm{sgn}(\overline{\Phi}_1(f)) = \mathrm{sgn}(\overline{\Phi}_1(f_{1'})) = 1$.

(2) Let $f \in O_{\mathbb{C}}(f_2) \cap \Omega_{\mathbb{R}}(2, 2)$. Note that $C_{\mathbb{C}}(f)$ is a parabola, hence $C_{\mathbb{R}}(f)$ is also a parabola. Moreover f has one cusp over $\mathbb{C}$ and since it is given by real equations the cusp must be real. Let $R_0 \in GA_{\mathbb{R}}(2, 2)$ be such that $C_{\mathbb{R}}(f \circ R_0) = \{2x^2 = y\}_{\mathbb{R}} = C_{\mathbb{R}}(f_2)$ and $f \circ R_0$ has the cusp at origin. Moreover let $L, R \in GA_{\mathbb{C}}(2, 2)$ be such that $f \circ R_0 = L \circ f_2 \circ R$. Note R is an automorphism of $C_{\mathbb{C}}(f \circ R_0) = \{2x^2 = y\}_{\mathbb{C}} = C_{\mathbb{C}}(f_2)$ and preserves the origin. Hence $R = (ax, a^2y)$ for some $a \in \mathbb{C}^*$. Let $L_1 = (a^2x, a^3y)$, note that we have $f_2 \circ R = L_1 \circ f_2$ and thus $f = L \circ L_1 \circ f_2 \circ R_0^{-1}$. Since the mapping $L \circ L_1$ maps $\mathbb{R}^2$ onto itself, we have $L \circ L_1 \in GA_{\mathbb{R}}(2, 2)$ and finally $f \in O_{\mathbb{R}}(f_2)$.

(3) Let $f \in O_{\mathbb{C}}(f_3) \cap \Omega_{\mathbb{R}}(2,2)$. Note that $C_{\mathbb{C}}(f)$ consists of two distinct intersecting lines, say $\{\alpha(x,y)=0\}$ and $\{\beta(x,y)=0\}$. Since $J(f)=\alpha\beta$ has real coefficients we may assume that α and β either have real or complex conjugate coefficients. However, the latter case can be excluded. Indeed, assume that α and β have complex conjugate coefficients. Since f has real coefficients $f(\overline{x},\overline{y}) = \overline{f(x,y)}$ and it follows that the components of the discriminant $\Delta(f)$ are also conjugate. This is a contradiction since the discriminant of mappings equivalent to f_3 consists of a line and a parabola.

So $C_{\mathbb{R}}(f)$ is a cross and $\Delta_{\mathbb{R}}(f)$ is a sum of a parabola and a tangent line. Let $L_0, R_0 \in GA_{\mathbb{R}}(2,2)$ be such that $C_{\mathbb{R}}(L_0 \circ f \circ R_0) = \{xy=0\}_{\mathbb{R}} = C_{\mathbb{R}}(f_3)$ and $\Delta_{\mathbb{R}}(L_0 \circ f \circ R_0) = \{y(y-x^2)=0\}_{\mathbb{R}} = \Delta_{\mathbb{R}}(f_3)$. Moreover let $L, R \in GA_{\mathbb{C}}(2,2)$ be such that $L_0 \circ f \circ R_0 = L \circ f_3 \circ R$. Note that R preserves the set $\{xy=0\}_{\mathbb{C}}$ and L preserves the set $\{y(y-x^2)=0\}_{\mathbb{C}}$, hence $L=(ax,a^2y)$ and $R=(bx,cy)$ for some $a,b,c \in \mathbb{C}^*$. Since $L \circ f_3 \circ R = (ab^2x^2+acy, a^2c^2y^2)$ is a real mapping we have $ac, ab^2 \in \mathbb{R}$. Moreover $f_3 = (b^2x, b^4y) \circ f_3 \circ (b^{-1}x, b^{-2}y)$, so $L_0 \circ f \circ R_0 = (ab^2x, a^2b^4y) \circ f_3 \circ (x, cb^{-2}y)$ and $f \in O_{\mathbb{R}}(f_3)$.

Note that $f \in O_{\mathbb{R}}(f_3)$ if and only if $\operatorname{sgn}(\overline{\Phi}_1(f)) = \operatorname{sgn}(\overline{\Phi}_1(f_3)) = -1$. Thus we have shown that for $f \in \Omega_{\mathbb{R}}(2,2)$ the conditions $\overline{\Phi}_1(f) \neq 0$, $\overline{\Phi}_2(f) = 0$ and $\operatorname{rank} \Psi_1(f) = 3$ imply $\overline{\Phi}_1(f) < 0$.

(4) Let $f \in O_{\mathbb{C}}(f_4) \cap \Omega_{\mathbb{R}}(2,2)$. Similarly as in (3) the set $C_{\mathbb{C}}(f)$ consists of two lines $\{\alpha(x,y)=0\}$ and $\{\beta(x,y)=0\}$ and we may assume that α and β either have real or complex conjugate coefficients.

If $C_{\mathbb{R}}(f)$ is a cross then we proceed similarly as in (3). We have $R_0 \in GA_{\mathbb{R}}(2,2)$ and $L, R \in GA_{\mathbb{C}}(2,2)$ such that $C_{\mathbb{R}}(f \circ R_0) = \{xy=0\}_{\mathbb{R}}$. Since R preserves the set $\{xy=0\}_{\mathbb{C}}$ and $f_4 = (a^{-2}x, b^{-2}y) \circ f_4 \circ (ax, by)$ we may assume that $R=(x,y)$ and deduce that $f \in O_{\mathbb{R}}(f_4)$.

Now let α and β have complex conjugate coefficients, we will show that $f \in O_{\mathbb{R}}(f_{4'})$, where $f_{4'} = (x^2-y^2, xy)$. Indeed, let $R_0 \in GA_{\mathbb{R}}(2,2)$ be such that $R_0(0,0)$ is the point of intersection of $\{\alpha(x,y)=0\}$ and $\{\beta(x,y)=0\}$ and $R_0(\{y-ix=0\}) = \{\alpha(x,y)=0\}$. It follows that $R_0(\{y+ix=0\}) = \{\beta(x,y)=0\}$ and consequently $C_{\mathbb{C}}(f \circ R_0) = \{x^2+y^2=0\}_{\mathbb{C}} = C_{\mathbb{C}}(f_{4'})$. Let $L, R \in GA_{\mathbb{C}}(2,2)$ be such that $f \circ R_0 = L \circ f_{4'} \circ R$. Note that R preserves $\{x^2+y^2=0\}_{\mathbb{C}}$ and we may assume that it does not switch the lines $\{y-ix=0\}$ and $\{y-ix=0\}$. So $R=(ax+by, -bx+ay)$ for some $a,b \in \mathbb{C}$ such that $a^2+b^2 \neq 0$. Let $L_1 = ((a^2-b^2)x + 4aby, -abx - (a^2-b^2)y)$, note that L_1 is an automorphism and $L_1 \circ f_{4'} = f_{4'} \circ R$. Thus $f \circ R_0 = L \circ L_1 \circ f_{4'}$, so $L \circ L_1 \in GA_{\mathbb{R}}(2,2)$ and $f \in O_{\mathbb{R}}(f_{4'})$.

As in (1) we obtain that $f \in O_{\mathbb{R}}(f_4)$ if and only if $\overline{\Phi}_1(f) < 0$, and $f \in O_{\mathbb{R}}(f_{4'})$ if and only if $\overline{\Phi}_1(f) > 0$.

(5) The description of orbits of f_5–f_{17} provided in Section 2 can be done in exactly the same way over $\mathbb{R}$. □

4. Final Remarks

At the end of this paper we state two conjectures:

Conjecture 1. *For every $d_1, d_2 > 0$ the set U of generic mappings in $\Omega(d_1,d_2)$ is an open affine subvariety of $\Omega(d_1,d_2)$. In particular every generic mapping is topologically stable, i.e. remains generic after a small deformation.*

Conjecture 2. *For every $d_1, d_2 > 0$ a mapping $f \in \Omega(d_1, d_2)$ is generic if $\Delta(f)$ contains only cusps and nodes and the number of cusps and nodes is maximal possible (see* [2]*).*

Note that for $d_1 = d_2 = 2$ these Conjectures are true, which follows from our paper.

References

[1] K. Aoki, H. Noguchi, On topological types of polynomial map germs of plane to plane, *Mem. School Sci. Eng. Waseda Univ.* 44, (1980), 133-156.

[2] M. Farnik, Z. Jelonek, M. Ruas, *Effective Whitney theorem for complex polynomial mappings of the plane*, arXiv:1503.00017v3 [math.AG], 2016.

[3] Z. Jelonek, *On finite regular and holomorphic mappings*, arXiv:1404.7466v4 [math.AG], 2015.

[4] Z. Jelonek, *A number of points in the set $\mathbb{C}^2 \setminus F(\mathbb{C}^2)$*, Bull. Polish Acad. Sci. Math. 47(3), (1999), 257-261.

[5] Z. Jelonek, *On semi-equivalence of generically-finite polynomial mappings*, Math. Z. 283(1), (2016), 133142.

[6] Z. Jelonek, K. Kurdyka, *Quantitative Generalized Bertini-Sard Theorem for smooth affine varieties*, Discrete and Computational Geometry 34, (2005), 659-678.

[7] T. K. Moe, *Rational Cuspidal Curves*, Department of Mathematics Faculty of Mathematics and Natural Sciences, University of Oslo, May 2008.

(M. Farnik) Jagiellonian University, Faculty of Mathematics and Computer Science, Łojasiewicza 6, 30-348 Kraków, Poland

E-mail address: michal.farnik@gmail.com

(Z. Jelonek) Instytut Matematyczny, Polska Akademia Nauk, Śniadeckich 8, 00-656 Warszawa, Poland

E-mail address: najelone@cyf-kr.edu.pl